

# Dimension spectra of random subfractals of self-similar fractals

Xiaoyang Gu\*    Jack H. Lutz†    Elvira Mayordomo‡ § ¶    Philippe Moser|| §

## Abstract

The dimension of a point  $x$  in Euclidean space (meaning the constructive Hausdorff dimension of the singleton set  $\{x\}$ ) is the *algorithmic information density* of  $x$ . Roughly speaking, this is the least real number  $\dim(x)$  such that  $r \times \dim(x)$  bits suffice to specify  $x$  on a general-purpose computer with arbitrarily high precision  $2^{-r}$ . The *dimension spectrum* of a set  $X$  in Euclidean space is the subset of  $[0, n]$  consisting of the dimensions of all points in  $X$ .

The dimensions of points have been shown to be geometrically meaningful (Lutz 2003, Hitchcock 2003), and the dimensions of points in self-similar fractals have been completely analyzed (Lutz and Mayordomo 2008). Here we begin the more challenging task of analyzing the dimensions of points in random fractals. We focus on fractals that are randomly selected subfractals of a given self-similar fractal. We formulate the specification of a point in such a subfractal as the outcome of an infinite two-player game between a *selector* that selects the subfractal and a *coder* that selects a point within the subfractal. Our selectors are algorithmically random with respect to various probability measures, so our selector-coder games are, from the coder's point of view, *games against nature*.

We determine the dimension spectra of a wide class of such randomly selected subfractals. We show that each such fractal has a dimension spectrum that is a closed interval whose endpoints can be computed or approximated from the parameters of the fractal. In general, the maximum of the spectrum is determined by the degree to which the coder can *reinforce* the randomness in the selector, while the minimum is determined by the degree to which the coder can *cancel* randomness in the selector. This constructive and destructive interference between the players' randomnesses is somewhat subtle, even in the simplest cases. Our proof techniques include van Lambalgen's theorem on independent random sequences, measure preserving transformations, an application of network flow theory, a Kolmogorov complexity lower bound argument, and a nonconstructive proof that this bound is tight.

---

\*Twitter, Inc. 1355 Market St, Suite 900, San Francisco, CA 94103, USA Email: xiaoyanggu@gmail.com

†Department of Computer Science, Iowa State University, Ames, IA 50011, USA. Email: lutz@cs.iastate.edu  
Research supported in part by National Science Foundation Grants 0652569 and 0728806, and by Spanish Government MEC Grant TIN 2005-08832-C03-02. Part of this work was done during a sabbatical at Caltech and the Isaac Newton Institute for Mathematical Sciences at the University of Cambridge.

‡Departamento de Informática e Ingeniería de Sistemas, Instituto de Investigación en Ingeniería de Aragón, Universidad de Zaragoza, 50018 Zaragoza, Spain. Email: elvira(at)unizar.es

§Research supported in part by Spanish Government MEC Grants TIN2005-08832-C03-02, TIN2008-06582-C03-02, and TIN2011-27479-C04-01.

¶Part of this author's research was performed during a visit at Iowa State University, supported by Spanish Government (Secretaría de Estado de Universidades e Investigación del Ministerio de Educación y Ciencia) grant for research stays PR2007-0368.

||Department of Computer Science, National University of Ireland, Maynooth. Maynooth, Co. Kildare. Ireland. Email: pmoser(at)cs.nuim.ie.

# 1 Introduction

Fractals are inherently information-theoretic objects. The dimension  $n$  of a Euclidean space  $\mathbb{R}^n$  is a measure of the amount of information (number of real numbers) that suffices to specify a point in  $\mathbb{R}^n$  in a natural way. Similarly, the fact that the Hausdorff dimension of the Cantor “middle-thirds” set  $C$  is  $\dim_H(C) = \log 2 / \log 3 \approx 0.63$  tells us that it only takes about 0.63 of a real number to specify a point in  $C$  in a natural way. That is, roughly  $(0.63)r$  bits suffice to specify the first  $r$  bits of a point in  $C$ . Intuitively, then, the Hausdorff (fractal) dimension  $\dim_H(C)$  is an upper bound on the “information densities” of points in the fractal  $C$ .

Of course some points in the Cantor set can be specified even more concisely. The theory of constructive dimension, a computability-theoretic extension of Hausdorff dimension developed in the present century [16], assigns each *individual point*  $x$  in a Euclidean space  $\mathbb{R}^n$  a *dimension*  $\dim(x) \in [0, n]$  that is a measure of its information density. This notion of dimension has been shown to be geometrically meaningful. For example, if  $X \subseteq \mathbb{R}^n$  is a “reasonably simple” set, in the sense that  $X$  is a union of  $\Pi_1^0$  (i.e., computably closed) sets, then

$$\dim_H(X) = \sup_{x \in X} \dim(x), \tag{1.1}$$

which is a nonclassical, pointwise characterization of the classical Hausdorff dimensions of such sets [16, 12].

The *self-similar fractals* form the best known and best understood class of fractals. (See Section 2.3 for a detailed review of self-similar fractals.) Each self-similar fractal  $F$  is given by an *iterated function system (IFS)*  $S = (S_1, \dots, S_{m-1})$  of *contracting similarities*  $S_i$ . A well-known theorem of Moran [20] states that

$$\dim_H(F) = \text{sdim}(F) \tag{1.2}$$

holds for every self-similar fractal  $F$ , where  $\text{sdim}(F)$  is the *similarity dimension* of  $F$ . Much of the importance of this theorem arises from the fact that  $\text{sdim}(F)$  is easy to compute from the contraction ratios  $c_0, \dots, c_{m-1}$  of the respective similarities  $S_1, \dots, S_{m-1}$ . That is, (1.2) gives an easy way to compute the Hausdorff dimensions of self-similar fractals.

The dimensions of points in computably self-similar fractals (those for which  $S_1, \dots, S_{m-1}$  are computable) have now been completely analyzed. If  $F$  is a self-similar fractal as above, then each point  $x \in F$  is naturally given by at least one *coding sequence*  $U \in \Sigma_m^\infty$ , where  $\Sigma_m = \{0, \dots, m-1\}$ . Intuitively,  $x$  is the result of a limiting process in which, at each stage  $t \in \mathbb{N}$ , we apply the contracting similarity  $S_{U[t]}$ . The main theorem of [18] says that, if  $F$  is computably self-similar, then, for each  $x \in F$  and each coding sequence  $U$  for  $x$ ,

$$\dim(x) = \text{sdim}(F) \dim^{\pi_S}(U), \tag{1.3}$$

where  $\dim^{\pi_S}(U) \in [0, 1]$  is the dimension of the sequence  $U \in \Sigma_m^\infty$  with respect to the *similarity probability measure*  $\pi_S$  on  $\Sigma_m$ , which arises from the IFS  $S$  in a natural manner. (This is a constructive version of Billingsley dimension [3] introduced in [18].)

This paper begins the more challenging task of analyzing the dimensions of points in random fractals. We focus on a particular class of random fractals, the *random subfractals* of self-similar fractals. For a concrete example, let  $F$  be the Sierpinski triangle. This is a well-known self-similar fractal. Intuitively, consider a *selector*  $\sigma$  that randomly chooses just two of the three top-level subtriangles of  $F$ , then randomly chooses just two subtriangles of each of these, etc., ultimately obtaining a subfractal  $F_\sigma$  of  $F$ . If  $\sigma$  is algorithmically random (with respect to some probability

distribution), then we call  $F_\sigma$  a *random subfractal* of  $F$ . An individual element  $x \in F_\sigma$  is specified by a *coder*  $T \in \{0, 1\}^\infty$  that tells us, at successive stages, *which* of the two selected subtriangles has  $x$  as an element. This interplay between  $\sigma$  and  $T$  is formalized in general terms as the *selector-coder game* in Section 3. Since our selectors are all random, our coders are playing games against nature.

The *dimension spectrum* of a set  $X \subseteq \mathbb{R}^n$  is the set  $\text{sp}(X) \subseteq [0, n]$  consisting of all  $\dim(x)$  for  $x \in X$ . Our objective is to determine the dimension spectrum  $\text{sp}(F_\sigma)$  of random subfractals of a given computably self-similar fractal  $F$ . By (1.3) and the fact that  $\dim^{\pi_S}(U)$  takes on all values in  $[0, 1]$ , we have

$$\text{sp}(F_\sigma) \subseteq \text{sp}(F) = [0, \text{sdim}(F)] \quad (1.4)$$

in any case.

Our main theorem, Theorem 4.1, concerns *similarity-random subfractals*  $F_\sigma$  of a given self-similar fractal  $F$ , i.e., subfractals of  $F$  specified by a selector  $\sigma$  that is algorithmically random with respect to a natural extension of the above-mentioned similarity probability measure  $\pi_S$  on  $\Sigma_m$ . This theorem says that each such  $\text{sp}(F_\sigma)$  is an interval containing  $\text{sdim}(F)$ , and it gives upper and lower bounds on the left endpoint of the interval  $\text{sp}(F_\sigma)$ . In the particular case where the contraction ratios  $c_0, \dots, c_{m-1}$  are all the same, these upper and lower bounds coincide, and our main theorem gives the exact dimension spectrum of  $F_\sigma$ .

Intuitively, the proof that  $\text{sdim}(F) \in \text{sp}(F_\sigma)$  is carried out by showing that the coder  $T$  can *reinforce* the randomness in  $\sigma$ , while the bounds on the left endpoint of  $\text{sp}(F_\sigma)$  quantify the degree to which the coder  $T$  can *cancel* some of the randomness in  $\sigma$ . This constructive and destructive interference between the players' randomnesses is somewhat subtle, and the proof of our main theorem reflects this, using van Lambalgen's theorem on independent random sequences, measure-preserving transformations, a Kolmogorov complexity lower bound argument, and a nonconstructive proof that this lower bound is nearly (and, in the single-contraction-ratio case, exactly) tight.

In Sections 5 and 6 we give results on the dimension spectra of subfractals of self-similar fractals that are random with respect to more general probability measures. Our proofs use the above methods, together with network flow theory and the divergence formula for randomness and dimension [17].

The randomness cancellation phenomena that play such a large role here have also arisen in other contexts, notably dimension spectra of random closed sets [2, 7] and of random translations of the Cantor set [8]. Our work is as much an investigation of these phenomena as it is an analysis of a particular class of fractals.

## 2 Preliminaries

### 2.1 Notation and terminology

Given a finite alphabet  $\Sigma$ , we write  $\Sigma^*$  for the set of all (finite) *strings* over  $\Sigma$  and  $\Sigma^\infty$  for the set of all (infinite) *sequences* over  $\Sigma$ . If  $\psi \in \Sigma^* \cup \Sigma^\infty$  and  $0 \leq i \leq j < |\psi|$ , where  $|\psi|$  is the length of  $\psi$ , then  $\psi[i]$  is the  $i$ th symbol in  $\psi$  (where  $\psi[0]$  is the leftmost symbol in  $\psi$ ), and  $\psi[i..j]$  is the string consisting of the  $i$ th through the  $j$ th symbols in  $\psi$ . If  $w \in \Sigma^*$  and  $\psi \in \Sigma^* \cup \Sigma^\infty$ , then  $w$  is a prefix of  $\psi$ , and we write  $w \sqsubseteq \psi$ , if there exists  $i \in \mathbb{N}$  such that  $w = \psi[0..i-1]$ . If  $A \subseteq \Sigma^*$  then  $A^{=n} = \{x \mid x \in A \wedge |x| = n\}$ .

For  $k \in \mathbb{N}$ ,  $\Sigma_k = \{0, \dots, k-1\}$ . Let  $s_0^{(k)}, s_1^{(k)}, s_2^{(k)}, \dots$  be the standard enumeration of  $\Sigma_k^*$ . For  $w \in \Sigma_k^*$ ,  $\text{index}^{(k)}(w)$  is the index of  $w$  in the standard enumeration of  $\Sigma_k^*$ , i.e.,  $s_{\text{index}^{(k)}(w)}^{(k)} = w$ .

For a set  $A$  and  $k \in \mathbb{N}$ ,  $[A]^k$  is the set of all  $k$ -element subsets of  $A$ . Given a set  $A \in [\Sigma_m]^k$ ,  $j \in \Sigma_k$ , we denote as  $A_j$  the  $j$ th element of  $A$  in alphabetical order.

For functions on Euclidean space, we use the computability notion formulated by Grzegorzczuk [11] and Lacombe [14] in the 1950's and expositied in the monographs by Pour-El and Richards [22], Ko [13], and Weihrauch [24] and in the recent survey paper by Braverman and Cook [5]. A function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is *computable* if there is an oracle Turing machine  $M$  with the following property. For all  $x \in \mathbb{R}^n$  and  $r \in \mathbb{N}$ , if  $M$  is given a function oracle  $\varphi_x : \mathbb{N} \rightarrow \mathbb{Q}^n$  such that, for all  $k \in \mathbb{N}$ ,  $|\varphi_x(k) - x| \leq 2^{-k}$ , then  $M$ , with oracle  $\varphi_x$  and input  $r$ , outputs a rational point  $M^{\varphi_x}(r) \in \mathbb{Q}^n$  such that  $|M^{\varphi_x}(r) - f(x)| \leq 2^{-r}$ .

For subsets of Euclidean space, we use the computability notion introduced by Brattka and Weihrauch [4] (see also [24, 5]). A set  $X \subseteq \mathbb{R}^n$  is computable if there is a computable function  $f_X : \mathbb{Q}^n \times \mathbb{N} \rightarrow \{0, 1\}$  that satisfies the following two conditions for all  $q \in \mathbb{Q}^n$  and  $r \in \mathbb{N}$ .

- (i) If there exists  $x \in X$  such that  $|x - q| \leq 2^{-r}$ , then  $f_X(q, r) = 1$ .
- (ii) If there is no  $x \in X$  such that  $|x - q| \leq 2^{1-r}$ , then  $f_X(q, r) = 0$ .

All logarithms in this paper are base-2. We use  $\mu$  for the uniform probability measures for Cantor spaces with all finite alphabet sizes.

## 2.2 Randomness and dimension

Recent advances in computability theory have yielded notions of dimension for single points in Euclidean spaces [16, 1, 18] (as opposed to the classical view where single points have dimension zero). These notions are robust in the sense that they admit several equivalent characterizations. We start with a description of constructive dimension on  $\Sigma_m^\infty$ . For the rest of this section let  $\pi$  denote a positive probability measure on  $\Sigma_m$ , extended by product to  $\Sigma_m^*$  (that is, extended by product to a probability measure on  $\Sigma_m^n$  for each  $n \in \mathbb{N}$ ).

**Definition.** Let  $s \in [0, \infty)$ . A  $\pi$ -*s-gale* is a function  $d : \Sigma_m^* \rightarrow [0, \infty)$  such that for all  $w \in \Sigma_m^*$ ,  $d(w)\pi^s(w) = \sum_{b \in \Sigma_m} d(wb)\pi^s(wb)$ . A  $\pi$ -1-gale is called a  $\pi$ -*martingale*. A  $\pi$ -*s-gale* is *constructive* (*lower semicomputable*) if there exists a computable function  $\hat{d} : \Sigma_m^* \times \mathbb{N} \rightarrow \mathbb{Q}$  such that

1. for all  $w, t$ ,  $\hat{d}(w, t) < d(w)$ , and
2. for all  $w$ ,  $\lim_{t \rightarrow \infty} \hat{d}(w, t) = d(w)$ .

An *optimal constructive  $\pi$ -martingale*  $d$  is constructive  $\pi$ -martingale such that for any constructive  $\pi$ -martingale  $d'$ , there is a constant  $c$  such that  $d(w) \geq cd'(w)$  for any  $w$ .

A  $\pi$ -*s-martingale succeeds* on  $T \in \Sigma_m^\infty$  if  $\limsup_{n \rightarrow \infty} d(T[0..n]) = \infty$ . A  $\pi$ -*s-martingale succeeds strongly* on  $T \in \Sigma_m^\infty$  if  $\liminf_{n \rightarrow \infty} d(T[0..n]) = \infty$ .

For a  $\pi$ -*s-gale*  $d$ , define its *success set* by  $S^\infty[d] = \{T \in \Sigma_m^\infty \mid \limsup_{n \rightarrow \infty} d(T[0..n]) = \infty\}$  and its *strong success set* by  $S_{\text{str}}^\infty[d] = \{T \in \Sigma_m^\infty \mid \liminf_{n \rightarrow \infty} d(T[0..n]) = \infty\}$ .

A sequence  $T \in \Sigma_m^\infty$  is  $\pi$ -*random* if no constructive  $\pi$ -martingale succeeds on it.

**Definition.** Let  $X \subseteq \Sigma_m^\infty$ . The (constructive) dimension of  $X$  relative to  $\pi$  is

$$\dim^\pi(X) = \inf\{s \in [0, \infty) \mid X \subseteq S^\infty[d] \text{ for some lower semicomputable } \pi\text{-}s\text{-gale } d\}.$$

For a sequence  $T \in \Sigma_m^\infty$  we write  $\dim^\pi(T)$  for  $\dim^\pi(\{T\})$ .

The constructive dimension of a sequence  $T$  characterizes the information density of the sequence. In fact, an alternative Kolmogorov complexity characterization of dimension was given in [19, 1, 18]. Fix a universal prefix free TM  $U$ . For any string  $x$ , the prefix-complexity of  $x$  denoted  $K(x)$ , is the size of the shortest binary program  $p$  such that  $U$  on input  $p$  produces  $x$ . The definition does not depend on the choice for  $U$  up to an additive constant.

**Definition.** Let  $w \in \Sigma_m^*$ . The *Shannon self-information* of  $w$  with respect to  $\pi$  is  $\mathcal{I}_\pi(w) = \log \frac{1}{\pi(w)} = \sum_{i=0}^{|w|-1} \log \frac{1}{\pi(w[i])}$ , where the logarithm is base-2 [6].

**Theorem 2.1** [19, 16, 1, 18] *Let  $T \in \Sigma_m^\infty$ . Then*

$$\dim^\pi(T) = \liminf_{j \rightarrow \infty} \frac{K(T[0..j-1])}{\mathcal{I}_\pi(T[0..j-1])}. \quad (2.1)$$

We now describe a similar dimension notion for single points in Euclidean spaces. For any  $x \in \mathbb{R}^n$  and  $r \in \mathbb{N}$ , consider the Kolmogorov complexity of  $x$  at precision  $r$  given by

$$K_r(x) = \min\{K(q) \mid q \in \mathbb{Q}^n \text{ and } |x - q| \leq 2^{-r}\}.$$

The dimension of points in  $\mathbb{R}^n$  is defined similarly to the Kolmogorov characterization in Theorem 2.1.

**Definition.** Let  $x \in \mathbb{R}^n$ . The *dimension* of  $x$  is

$$\dim(x) = \liminf_{r \rightarrow \infty} \frac{K_r(x)}{r}. \quad (2.2)$$

**Definition.** Let  $X \subseteq \mathbb{R}^n$ . The (constructive) dimension of  $X$  is

$$\dim(X) = \sup_{x \in X} \dim(x).$$

### 2.3 Self-similar fractals

Many famous fractals are made up of parts that are resized copies of the whole set. As an example, the middle third Cantor set is the union of two shrunken versions of the whole set. Such fractals are called self-similar.

Self-similar fractals are formally defined as the unique invariant set under a family of contracting similarities  $S_0, \dots, S_{m-1}$ , where for every  $i \leq m-1$  and  $x, y \in \mathbb{R}^n$ ,  $S_i : D \rightarrow D$  (where  $D \subset \mathbb{R}^n$  is closed) and  $|S_i(x) - S_i(y)| = c_i|x - y|$ , where  $c_i \in (0, 1)$ .

With the example of the middle third Cantor set  $F$ , letting  $S_0(x) = 1/3x$  and  $S_1(x) = 1/3x + 2/3$  ( $S_0, S_1 : \mathbb{R} \rightarrow \mathbb{R}$ ) the invariance property can be expressed as  $F = S_0(F) \cup S_1(F)$ , where  $S_0(F)$  (resp.  $S_1(F)$ ) is the resized copy of  $F$  placed on the left (resp. right).

Formally a finite sequence  $S = (S_0, \dots, S_{m-1})$  of two or more contracting similarities on a nonempty, closed set  $D \subseteq \mathbb{R}^n$ , is called an *iterated function system (IFS)* [10]. We call  $D$  the *domain* of  $S$ , writing  $D = \text{dom}(S)$ . Each  $S_i$  has a contraction ratio  $c_i \in (0, 1)$ . Let  $\mathcal{K}(D)$  be the set of compact subsets of  $D$ , and  $\mathcal{K}(S) = \mathcal{K}(\text{dom}(S))$ .  $S$  induces a function  $S : \mathcal{K}(S) \rightarrow \mathcal{K}(S)$  defined by  $S(A) = \cup_{i=0}^{m-1} S_i(A)$ .

Going back to the example of the middle third Cantor set, using the alphabet  $\Sigma = \{0, 1\}$  to refer to the contracting similarities  $S_0$  and  $S_1$ , then each point  $P$  in  $F$  can be specified by

an infinite sequence  $T \in \Sigma^\infty$ , that codes for the infinite sequence of similarities  $S_{T[0]}, S_{T[1]}, \dots$  that when applied successively to  $A = [0, 1]$ , yield  $P$ . Denoting  $P$  by  $S(T)$ , The middle third Cantor set can be expressed as the set of points encoded by all infinite sequences  $T \in \Sigma^\infty$ , i.e.  $F(S) = \{S(T) \mid T \in \Sigma^\infty\}$ .

The general case is similar: Let  $S = (S_0, \dots, S_{m-1})$  be an IFS. Let  $A \in \mathcal{K}(S)$  be such that  $S(A) \subseteq A$ . Consider the function  $S_A : \Sigma_m^* \rightarrow \mathcal{K}(S)$  defined by the following induction.  $S_A(\lambda) = A$  and  $S_A(wi) = S_i(S_A(w))$  for every  $w \in \Sigma_m^*$ ,  $i \in \Sigma_m$ . Because all the contraction ratios  $c_i$  are smaller than 1, it is easy to see that for each sequence  $T \in \Sigma_m^\infty$  there is a unique point  $S_A(T) \in \mathbb{R}^n$  such that  $\bigcap_{w \sqsubseteq T} S_A(w) = \{S_A(T)\}$ . We call  $T$  a *coding sequence* for the point  $S_A(T)$ . It is well known [10] that the function  $S_A : \Sigma_m^\infty \rightarrow \mathbb{R}^n$  does not depend on the choice of  $A$  i.e., for any  $A, B \in \mathcal{K}(D)$  such that  $S(A) \subseteq A$  and  $S(B) \subseteq B$ , we have  $S_A = S_B$ . Thus for any IFS  $S$ , the function  $S : \Sigma_m^\infty \rightarrow \mathbb{R}^n$  obtained by letting  $S = S_A$  for some set  $A \in \mathcal{K}(D)$  such that  $S(A) \subseteq A$  is well defined. The *attractor* of the IFS  $S$  is given by  $F(S) = \{S(T) \mid T \in \Sigma_m^\infty\}$ .

Because the sets  $S_0(D), \dots, S_{m-1}(D)$  need not be disjoint, a point in  $F(S)$  can admit more than one coding sequence  $T$ .

An attractor  $F$  of an IFS  $S$  is a *self-similar fractal* if the sets  $S_0(D), \dots, S_{m-1}(D)$  are “almost” disjoint i.e., if  $F$  satisfies the following *open set condition*.

**Definition.** An IFS  $S = (S_0, \dots, S_{m-1})$  with domain  $D$  satisfies the *open set condition* if there exists a nonempty, bounded, open set  $G \subseteq D$  such that  $S_0(G), \dots, S_{m-1}(G)$  are disjoint subsets of  $G$ .

It is a classical result of Moran [20] and Falconer [9] that for any self similar fractal  $F(S)$ , the box, packing and Hausdorff dimension all coincide and are equal to the unique solution  $\text{sdim}(F)$  of equation  $\sum_{i=0}^{m-1} c_i^{\text{sdim}(F)} = 1$ , where the  $c_i$ 's are the contraction ratios of  $S$ .

Iterated function systems induce probability measures on alphabets in the following manner.

**Definition.** The *similarity probability measure* of an IFS  $S = (S_0, \dots, S_{m-1})$  with contraction ratios  $c_0, \dots, c_{m-1}$  is the probability measure  $\pi_S$  on the alphabet  $\Sigma_m$  defined by

$$\pi_S(i) = c_i^{\text{sdim}(F)}$$

for all  $i \in \Sigma_m$ .

In this paper, we are interested in computable IFSs. Here is a definition.

**Definition.** An IFS  $S = (S_0, \dots, S_{m-1})$  is computable if  $\text{dom}(S)$  is a computable set and the functions  $S_0, \dots, S_{m-1}$  are computable.

**Definition.** A *computably self-similar fractal* is a set  $F \subseteq \mathbb{R}^n$  that is the attractor of an IFS that is computable and satisfies the open set condition.

The following theorem is our starting point.

**Theorem 2.2** [18]. *If  $F \subseteq \mathbb{R}^n$  is a computably self-similar fractal given by a computable IFS  $S$ , then, for all points  $x \in F$  and all coding sequences  $U$  of  $x$ ,*

$$\dim(x) = \text{sdim}(F) \dim^{\pi_S}(U).$$

### 3 The selector-coder game

The random fractals that we consider in this paper are randomly selected subfractals of a given computably self-similar fractal. This section explains this random selection process in terms of a

two-player game. As elaborated below, each outcome of this game codifies a point in the random fractal and the power of the players in the game captures different possibilities for the fractal randomness.

Let  $n$ ,  $m$ , and  $k$  be integers with  $n \geq 1$  and  $m \geq k \geq 2$ . Let  $F = F(S) \subseteq \mathbb{R}^n$  be a self-similar fractal given by a computable IFS  $S = (S_0, \dots, S_{m-1})$  satisfying the open set condition. Recall that

$$F = \{S(U) \mid U \in \Sigma_m^\infty\},$$

i.e., each point  $x \in F$  is of the form  $x = S(U)$  for some coding sequence  $U \in \Sigma_m^\infty$ . We are interested in certain randomly selected subfractals of the fractal  $F$ . It is easiest to specify such a subfractal by saying *which* coding sequences  $U \in \Sigma_m^\infty$  give rise to points  $S(U)$  in the subfractal.

Intuitively, our subfractal consists of points  $S(U)$  for which  $U$  is the outcome of a game played by a *selector* and a *coder*. During each round  $t = 0, 1, \dots$  of this game, the  $t$ th symbol  $U[t] \in \Sigma_m$  of  $U$  is determined by the following choices.

- (i) The selector chooses a  $k$ -element subset  $A$  of  $\Sigma_m$ .
- (ii) The coder chooses an element  $i$  of  $\Sigma_k$ .

The symbol  $U[t]$  is then the  $i$ th element of  $A$ .

More formally, recall that  $[\Sigma_m]^k$  is the set of all  $k$ -element subsets of  $\Sigma_m$ . Given a set  $A \in [\Sigma_m]^k$  and an index  $i \in \Sigma_k$ , we let  $A_i$  denote the  $i$ th element of  $A$  in the standard ordering of  $\Sigma_m$ . Thus  $A = \{A_0, \dots, A_{k-1}\}$  and  $A_0 < \dots < A_{k-1}$ .

The following definition allows the selector's choice of the set  $A$  to depend upon the coder's earlier choices of symbols in  $\Sigma_k$ .

**Definition.** An  $\binom{m}{k}$ -*selector* (or, when  $m$  and  $k$  are clear from the context, a *selector*) is a function

$$\sigma : \Sigma_k^* \rightarrow [\Sigma_m]^k.$$

We write  $\text{SEL}\binom{m}{k}$  for the set of all  $\binom{m}{k}$ -selectors.

For the purpose defining random subfractals, it does not really matter *how* the coder makes its choices, i.e., we can identify the coder with the sequence of choices that it makes.

**Definition.** A *coder* is a sequence  $T \in \Sigma_k^\infty$ .

Once a selector and a coder have been chosen, the outcome of the selector-coder game is determined.

**Definition.** Let  $\sigma \in \text{SEL}\binom{m}{k}$  be a selector, and let  $T \in \Sigma_k^\infty$  be a coder. The *outcome* of (the selector-coder game played between)  $\sigma$  and  $T$  is the sequence  $\sigma * T \in \Sigma_m^\infty$  defined by

$$(\sigma * T)[t] = \sigma(T[0..t-1])_{T[t]}$$

for all  $t \in \mathbb{N}$ .

Our intent, captured in the next definition, is for each selector  $\sigma$  to specify (select) the subfractal  $F_\sigma$  of  $F$  consisting of all points  $S(U)$  for which  $U$  is an outcome of playing  $\sigma$  against some coder.

**Definition.** For each selector  $\sigma \in \text{SEL}\binom{m}{k}$ , the *subfractal* of  $F$  selected by  $\sigma$  is the set

$$F_\sigma = \{S(\sigma * T) \mid T \in \Sigma_k^\infty\}.$$

The following observation, which follows immediately from Theorem 2.2, reduces our study of the dimensions of points in  $F_\sigma$  to a study of the dimensions of sequences of the form  $\sigma * T$ .

**Observation 3.1** For each point  $x = S(\sigma * T) \in F_\sigma$  we have

$$\dim(x) = \text{sdim}(F) \dim^{\pi_S}(\sigma * T).$$

Our interest here is in randomly selected subfractals of  $F$ , by which we mean subfractals  $F_\sigma$  of  $F$  for which the selector  $\sigma$  is random with respect to some probability measure. That is, we are interested in the case where the coder is playing a “game against nature”. To make this idea precise, we identify each selector  $\sigma : \Sigma_k^* \rightarrow [\Sigma_m]^k$  with its *characteristic* sequence  $\chi_\sigma \in ([\Sigma_m]^k)^\infty$  defined by

$$\chi_\sigma[i] = \sigma(s_i^{(k)})$$

for all  $i \in \mathbb{N}$ . Note that  $\chi_\sigma$  is an infinite sequence over the  $\binom{m}{k}$ -element *alphabet*  $[\Sigma_m]^k$ .

We now have two ways to visualize a selector  $\sigma$ . The original definition suggests that we regard a selector  $\sigma : \Sigma_k^* \rightarrow [\Sigma_m]^k$  as a *labeled tree* in which this underlying tree is  $\Sigma_k^*$  (i.e., the root is  $\lambda$ , and each vertex  $w$  has the left-to-right children  $w0, w1, \dots, w(k-1)$ ) and each vertex  $w$  has the set  $\sigma(w) \in [\Sigma_m]^k$  as its label. The identification of  $\sigma$  with its characteristic sequence  $\chi_\sigma \in ([\Sigma_m]^k)^\infty$  suggests that we regard  $\sigma$  as a *breadthfirst traversal* of this labeled tree. As we shall see, these are both useful perspectives.

We often analyze selectors, coders, and their outcomes in terms of finite “initial segments”. To this end, we define a string  $u \in ([\Sigma_m]^k)^*$  to be a *prefix* of a selector  $\sigma$ , and we write  $u \sqsubseteq \sigma$ , if  $u \sqsubseteq \chi_\sigma$ .

We next define a “finite prefix version” of the outcome operation  $(\sigma, T) \mapsto \sigma * T$ . This takes a bit of care, because  $\sigma * T$  only depends on those values  $\sigma(v)$  of  $\sigma$  for which  $v \sqsubseteq T$ .

**Definition.** Let  $u \in ([\Sigma_m]^k)^*$  and  $v \in \Sigma_k^*$ . Let  $v'$  be the longest prefix of  $v$  such that  $j < |u|$  for every proper prefix  $s_j^{(k)} \sqsubsetneq v'$ . Then the *result* of  $u$  and  $v$  is the string  $u * v \in \Sigma_m^{|v'|}$  defined by

$$(u * v)[i] = u[\text{index}^{(k)}(v[0..i-1])_{v[i]}]$$

for all  $0 \leq i < |v'|$ , recall that  $\text{index}^{(k)}(w)$  is the index of  $w$  in the standard enumeration of  $\Sigma_k^*$ , i.e.,  $s_{\text{index}^{(k)}(w)}^{(k)} = w$ .

**Observation 3.2** If  $\sigma \in \text{SEL}(\binom{m}{k})$ ,  $T \in \Sigma_k^\infty$ ,  $u \sqsubseteq \sigma$ , and  $v \sqsubseteq T$ , then  $u * v$  is the longest prefix of  $\sigma * T$  that is determined by  $u$  and  $v$ .

We now define what it means for the selector  $\sigma$  to be random with respect to a given probability measure.

**Definition.** Let  $\gamma \in \Delta([\Sigma_m]^k)$ , i.e.,  $\gamma$  is a probability measure on the discrete sample space  $[\Sigma_m]^k$ . A selector  $\sigma \in \text{SEL}(\binom{m}{k})$  is *random* with respect to  $\gamma$  (or, more simply,  $\gamma$ -*random*) if its characteristic sequence  $\chi_\sigma$  is  $\gamma$ -random.

This paper is concerned with the following type of fractal.

**Definition.** Let  $\gamma \in \Delta([\Sigma_m]^k)$ . A  $\gamma$ -*random subfractal* of  $F$  is a set  $F_\sigma$ , where  $\sigma$  is a  $\gamma$ -random selector.

The following well-known Kolmogorov complexity characterization of  $\gamma$ -randomness is useful. The proof is a generalization of Theorem 3.6.1 in [15] using Theorem 3.3.1 in [15].

**Theorem 3.3** A selector  $\sigma \in \text{SEL}(\binom{m}{k})$  is random with respect to a computable probability measure  $\gamma \in \Delta([\Sigma_m]^k)$  if and only if every sufficiently long prefix  $u \sqsubseteq \sigma$  satisfies

$$K(u) \geq \mathcal{I}_\gamma(u) - c_\sigma.$$



## 4 Similarity-random subfractals

This is the main section of the paper. We investigate the dimension spectra of a natural class of random subfractals of a self-similar fractal. This class is somewhat restrictive, but it exhibits several subtleties of the interactions between randomness and dimension.

As before, let  $n$ ,  $m$ , and  $k$  be integers with  $n \geq 1$  and  $m \geq k \geq 2$ . Let  $F = F(S) \subseteq \mathbb{R}^n$  be a computably self-similar fractal given by a computable IFS  $S = (S_0, \dots, S_{m-1})$  satisfying the open set condition.

**Definition.** The *similarity probability measure* induced by  $S$  (equivalently, by  $F$ ) on  $[\Sigma_m]^k$  is the probability measure  $\hat{\pi}_S \in \Delta([\Sigma_m]^k)$  given by

$$\hat{\pi}_S(A) = \frac{\pi_S(A)}{\binom{m-1}{k-1}}$$

for each  $A \in [\Sigma_m]^k$ . Here  $\pi_s$  is the similarity probability measure on  $\Sigma_m$  defined in Section 2, and we write  $\pi_S(A) = \sum_{i \in A} \pi_S(i)$ .

It is routine to verify that  $\sum_{A \in [\Sigma_m]^k} \hat{\pi}_S(A) = 1$ , whence  $\hat{\pi}_S \in \Delta([\Sigma_m]^k)$ . It should also be noted that, if the similarities  $S_0, \dots, S_{m-1}$  all have the same contraction ratio, then  $\pi_s$  is the uniform probability measure on  $\Sigma_m$ , and  $\hat{\pi}_S$  is the uniform probability measure on  $[\Sigma_m]^k$ .

**Definition.**

1. A selector  $\sigma \in \text{SEL}(\binom{m}{k})$  is *similarity random* if it is  $\hat{\pi}_S$ -random.
2. A *similarity random subfractal* of  $F$  is a subfractal  $F_\sigma$  of  $F$  (as defined in Section 3), where  $\sigma$  is a similarity random selector.

Our objective is to prove the following result that analyzes the dimension spectrum of similarity-random subfractals, as explained informally in the introduction. In the particular case where the contraction ratios  $c_0, \dots, c_{m-1}$  are all the same, these upper and lower bounds coincide, and our main theorem gives the exact dimension spectrum of  $F_\sigma$ .

**Theorem 4.1** (main theorem) *For every similarity random subfractal  $F_\sigma$  of  $F$ , the dimension spectrum  $\text{sp}(F_\sigma)$  is an interval satisfying*

$$[s^* \frac{\log(m-1) - \log(k-1 + A(m-k))}{\log \frac{1}{a}}, s^*] \subseteq \text{sp}(F_\sigma) \subseteq [s^* \frac{\log m - \log k}{\log \frac{1}{A}}, s^*],$$

where  $s^* = \text{sdim}(F)$ ,  $a = \min\{\pi_S(i) \mid i \in \Sigma_m\}$ , and  $A = \max\{\pi_S(i) \mid i \in \Sigma_m\}$ .

*In particular, if all the contraction ratios of  $F$  have the same value  $c$ , then every similarity-random (i.e., uniformly random) subfractal  $F_\sigma$  of  $F$  has dimension spectrum*

$$\text{sp}(F_\sigma) = [s^*(1 - \frac{\log k}{\log m}), s^*],$$

where  $s^* = \text{sdim}(F) = (\log m)/(\log \frac{1}{c})$ .

**Example 4.2** *Let  $F$  be the standard Sierpinski triangle. This is given by an IFS  $S = (S_0, S_1, S_2)$  in which all three contraction ratios are  $c = \frac{1}{2}$ , so  $\dim_{\text{H}}(F) = \text{sdim}(F) = \log 3$ . If  $\sigma$  is a uniformly random selector that chooses two of the contractions  $S_0, S_1, S_2$  at every stage, then Theorem 4.1 says that the resulting random subfractal  $F_\sigma$  of  $F$  has dimension spectrum*

$$\text{sp}(F_\sigma) = [(\log 3) - 1, \log 3] \approx [0.585, 1.585].$$

We now turn to the proof of Theorem 4.1. Let  $J$  be the set of all possible values of  $\dim^{\pi_S}(\sigma * T)$ . By Observation 3.1 it suffices to prove the following three things.

- $J$  is an interval. (4.1)

- $1 \in J$  (4.2)

- $\frac{\log m - \log k}{\log \frac{1}{a}} \leq \inf J \leq \frac{\log m - \log k}{\log \frac{1}{A}}$  (4.3)

To see that (4.1) holds, let  $\alpha_1 < \beta < \alpha_2$ , with  $\alpha_1, \alpha_2 \in J$ . Then there exist  $T_1, T_2 \in \Sigma_k^\infty$  such that each  $\dim^{\pi_S}(\sigma * T_i) = \alpha_i$ . It follows by (2.1) that

$$\liminf_{j \rightarrow \infty} \frac{K((\sigma * T_i)[0..j-1])}{\mathcal{I}_{\pi_S}((\sigma * T_i)[0..j-1])} = \alpha_i$$

holds for  $i = 1, 2$ . It suffices to show that there is a sequence  $T \in \Sigma_k^\infty$  such that

$$\lim_{j \rightarrow \infty} \frac{k_j}{i_j} = \beta,$$

where

$$k_j = K((\sigma * T)[0..j-1])$$

and

$$i_j = \mathcal{I}_{\pi_S}((\sigma * T)[0..j-1])$$

for each  $j \in \mathbb{N}$ . We let

$$T = w_{10} w_{20} w_{11} w_{21} w_{12} w_{22} \dots,$$

where each  $w_{1k}$  is a prefix of  $T_1$  just long enough that

$$\frac{k_j}{i_j} < \beta,$$

where  $j = |w_{10} w_{20} \dots w_{1k}|$ , and each  $w_{2k}$  is a prefix of  $T_2$  just long enough that

$$\frac{k_j}{i_j} > \beta,$$

where  $j = |w_{10} w_{20} \dots w_{1k} w_{2k}|$ . These prefixes exist because  $\alpha_1 < \beta < \alpha_2$ , and  $\frac{k_j}{i_j}$  converges to  $\beta$ , because each symbol in  $T$  determines a corresponding symbol of  $\sigma * T$  and

$$\begin{aligned} \left| \frac{k_{j+1}}{i_{j+1}} - \frac{k_j}{i_j} \right| &\leq \frac{i_j |k_{j+1} - k_j| + k_j |i_{j+1} - i_j|}{i_j^2} \\ &= O\left(\frac{j \cdot 1 + j \cdot 1}{j^2}\right) \\ &= o(1) \end{aligned}$$

as  $j \rightarrow \infty$ . We now turn to the more interesting components of the proof of Theorem 4.1.

The following lemma and theorem establish that (4.2) holds.

**Lemma 4.3** *There is a distribution  $\gamma \in \Delta(\Sigma_k)$  such that the outcome operation  $(\sigma, T) \mapsto \sigma * T$  is measure-preserving when using distributions  $\pi_S \in \Delta(\Sigma_m)$  and  $\hat{\pi}_S \in \Delta([\Sigma_m]^k)$ .*

**Proof.** We explicitly obtain  $\gamma$  by solving the following system of equations, for  $i \in \Sigma_m$ ,

$$\pi_S(i) = \sum_{j \in \Sigma_k} \sum_{A \in [\Sigma_m]^k, A_j = i} \hat{\pi}_S(A) \gamma(j).$$

We use the notation  $\hat{\pi}_S(0) = \sum_{0 \in B} \hat{\pi}_S(B)$ .

Notice that  $\gamma(0)$  can be obtained from the first equation

$$\pi_S(0) = \hat{\pi}_S(0) \gamma(0),$$

then  $\gamma(1)$  can be obtained from

$$\pi_S(1) = \sum_{1 \in A, 0 \notin A} \hat{\pi}_S(A) \gamma(0) + \sum_{0, 1 \in A} \hat{\pi}_S(A) \gamma(1),$$

and so on. □

**Theorem 4.4** *Let  $\gamma$  be the distribution given by the previous lemma. If a coder  $T$  is  $\gamma$ -random relative to a similarity-random selector  $\sigma$ , then the coding sequence  $\sigma * T$  is  $\pi_S$ -random, so  $\dim^{\pi_S}(\sigma * T) = 1$ .*

**Proof.** The proof of this result uses the ideas in van Lambalgen's theorem [23] and the fact that the outcome operation  $(\sigma, T) \mapsto \sigma * T$  is measure-preserving (Lemma 4.3).

We prove this theorem by proving that if  $\sigma$  is  $\hat{\pi}_S$ -random and  $\sigma * T$  is not  $\pi_S$ -random, then  $T$  is not  $\gamma$ -random relative to  $\sigma$ .

Let  $d : \Sigma_m^\infty \rightarrow [0, \infty]$  be an optimal constructive  $\pi_S$ -martingale on  $\Sigma_m^\infty$ . Without loss of generality, assume that  $d(\lambda) < 1$ .

For each  $n \in \mathbb{N}$ , let

$$Z_n = \{U \in \Sigma_m^\infty \mid (\exists u \sqsubseteq U) d(u) > 2^{2n}\}.$$

It is clear that  $\{Z_n\}_{n \in \mathbb{N}}$  is uniformly computably enumerable (or  $\Sigma_1^0$ ). Since for every prefix-free  $A \subseteq \Sigma_m^*$  the (generalized) Kraft's inequality implies that  $\sum_{u \in A} d(u) \pi_S(u) \leq d(\lambda)$ , we have that  $\pi_S(Z_n) < 2^{-2n}$ .

For each  $\sigma' \in \text{SEL}(\binom{m}{k})$  and each  $n \in \mathbb{N}$ , let

$$Y_n^{\sigma'} = \{T' \in \Sigma_k^\infty \mid (\exists u \sqsubseteq \sigma' * T') d(u) > 2^{2n}\}.$$

Let

$$X_n = \left\{ \sigma' \in \text{SEL}(\binom{m}{k}) \mid \gamma(Y_n^{\sigma'}) > 2^{-n} \right\}.$$

It is clear that  $\{X_n\}_{n \in \mathbb{N}}$  is  $\Sigma_1^0$ . Then since  $*$  is measure preserving,  $\pi_S(Z_n) \geq \hat{\pi}_S(X_n) \cdot \gamma(Y_n^{\sigma'})$  and

$$\hat{\pi}_S(X_n) \cdot 2^{-n} \leq \pi_S(Z_n) < 2^{-2n}.$$

Therefore,  $\hat{\pi}_S(X_n) < 2^{-n}$ .

Since  $\sigma$  is  $\hat{\pi}_S$ -random,  $|\{n \in \mathbb{N} \mid \sigma \in X_n\}| < \infty$  and thus there exists  $n_0 \in \mathbb{N}$  such that for all  $n_0 \leq n \in \mathbb{N}$ ,

$$\gamma(Y_n^\sigma) \leq 2^{-n}.$$

Note that  $\{Y_n^\sigma\}_{n \in \mathbb{N}}$  is uniformly  $\Sigma_1^0$  in  $\sigma$ . Since  $\sigma * T$  is not  $\pi_S$ -random,  $T \in \bigcap_{n \in \mathbb{N}} Y_n^\sigma$ , which is a  $\sigma$ -effective measure 0 set. Therefore  $T$  is not  $\gamma$ -random relative to  $\sigma$ .  $\square$

We now turn to (4.3), which concerns the left endpoint of the interval  $J$ . The question is now how small the coder  $T$  can force the dimension  $\dim^{\pi_S}(\sigma * T)$  to be. More intuitively, how much of the randomness in  $\sigma$  can the coder “cancel”? The following theorem places an upper bound on the amount of such cancellation and thereby establishes the left-hand inequality in (4.3).

**Theorem 4.5** *If  $\sigma$  is a similarity-random selector, then for every coder  $T$ ,*

$$\dim^{\pi_S}(\sigma * T) \geq \frac{\log(m-1) - \log(k-1 + A(m-k))}{-\log a},$$

where  $a = \min\{\pi_S(i) \mid i \in \Sigma_m\}$ ,  $A = \max\{\pi_S(i) \mid i \in \Sigma_m\}$ .

The proof is a Kolmogorov complexity argument. Roughly, if a prefix  $w$  of  $\sigma * T$  can be compressed to  $(\frac{\log(m-1) - \log(k-1 + A(m-k))}{-\log a} - \epsilon)\mathcal{I}_{\pi_S}(w)$  bits, then  $\sigma$  is “somewhat” compressible, hence nonrandom by Theorem 3.3. The proof uses the following lemma about Kolmogorov complexity.

**Lemma 4.6** *There is a constant  $c \in \mathbb{N}$  with the following property. For every selector  $\sigma$ , every coder  $T$ , and every prefix  $w \sqsubseteq \sigma * T$ , if  $u_w$  is the  $(k^{|w|} - 1)/(k-1)$ -symbol prefix of  $\sigma$  (i.e., the prefix of  $\sigma$  that determines all labels at depth less than  $|w|$  when  $\sigma$  is viewed as a labeled tree), then*

$$K(u_w) \leq \mathcal{I}_{\hat{\pi}_S}(u_w) + K(w) - |w|(\log(m-1) - \log(k-1 + A(m-k))) + c. \quad (4.4)$$

**Proof.** In this proof we show that  $w$  possesses a significant amount of information about a prefix  $u_w$  of  $\sigma$ , and compressing  $w$  allows us to compress  $u_w$ .

(Recall that for a set  $A$  and  $k \in \mathbb{N}$ ,  $[A]^k$  is the set of all  $k$ -element subsets of  $A$ .)

We use the following notation. For  $i \in \Sigma_m$  we refer to  $\hat{\pi}_S(i)$  as the sum of  $\hat{\pi}_S$  for the sets containing  $i$ , that is

$$\hat{\pi}_S(i) = \sum_{i \in B} \hat{\pi}_S(B).$$

We abuse notation and for  $w \in \Sigma_m^*$  we use

$$\mathcal{I}_{\hat{\pi}_S}(w) = \sum_{j=0}^{|w|-1} \log \frac{1}{\hat{\pi}_S(w[j])}$$

From the definition of  $\hat{\pi}_S$  we have that

$$\hat{\pi}_S(i) = \frac{k-1}{m-1} + \pi_S(i) \frac{m-k}{m-1}.$$

We will prove that there is a constant  $c \in \mathbb{N}$  with the following property. For every selector  $\sigma$ , every coder  $T$ , and every prefix  $w \sqsubseteq \sigma * T$ , if  $u_w$  is the  $(k^{|w|} - 1)/(k-1)$ -symbol prefix of  $\sigma$ , then

$$K(u_w) \leq \mathcal{I}_{\hat{\pi}_S}(u_w) + K(w) - \mathcal{I}_{\hat{\pi}_S}(w) + c. \quad (4.5)$$

Notice that equation 4.5 implies the lemma.

Let us describe informally our proof of equation 4.5. We are going to code  $u_w$  by using  $\hat{\pi}_S$ -Shannon-Fano coding on strings in  $([\Sigma_m]^k)^{(k^{|w|}-1)/(k-1)}$  (that is, strings of length  $|u_w|$  over the

alphabet  $[\Sigma_m]^k$  will be coded according to their  $\hat{\pi}_S$ -probability). In fact we will not code the whole  $u_w$  but only the part of it not in the path corresponding to  $T$ . For the part in the  $T$  path we will first code  $w$  and then code the rest of the path with the help of  $w$ .

Let  $v$  be the prefix of  $T$  with  $|v| = |w|$ . Define  $y \in ([\Sigma_m]^{k-1})^{|w|}$  by

$$y[i] = \sigma(v[0..i-1]) - \{w[i]\}$$

for all  $0 \leq i < |w|$ . Let  $z$  be the string obtained from  $u_w$  by deleting all the symbols  $u_w[\text{index}^{(k)}(v')]$  for which  $\lambda \neq v' \sqsubseteq v$ . (Note that  $u_w[\text{index}^{(k)}(v')] = \sigma(v')$ ). Intuitively,  $z$  is  $u_w$  without the path corresponding to  $T$  and  $y$  is the path corresponding to  $T$  but removing an element ( $w[i]$ ) from each set.

We will use two different Shannon-Fano codes. In the first one, given  $w$ ,  $k$  and  $m$ , we code each string  $p \in ([\Sigma_m]^k)^{\frac{k|w|-1}{k-1}-|w|}$  with  $\lceil -\sum_{j=0}^{|p|-1} \log(\hat{\pi}_S(p[j])) \rceil$  bits using Shannon-Fano code with probability  $\gamma(p) = \prod_{j=0}^{|p|-1} \hat{\pi}_S(p[j])$ .

In the second code, given  $w$ ,  $k$  and  $m$ , we code each string  $q \in ([\Sigma_m]^{k-1})^{|w|}$  satisfying  $w[i] \notin q[i]$  for all  $0 \leq i < |w|$  with  $\lceil \sum_{j=0}^{|q|-1} (-\log(\hat{\pi}_S(q[j] \cup \{w[j]\})) + \log(\hat{\pi}_S(w[j]))) \rceil$  bits using Shannon-Fano code with probability  $\nu(q) = \prod_{j=0}^{|q|-1} \hat{\pi}_S(q[j] \cup \{w[j]\}) / \hat{\pi}_S(w[j])$ .

Then  $u_w$  can be coded by first coding  $w$ ,  $k$  and  $m$ , then coding  $z$  with the first code and finally coding  $y$  with the second code.

Therefore,

$$K(u_w) \leq K(w) + K(k, m) + \lceil -\sum_{j=0}^{|z|-1} \log(\hat{\pi}_S(z[j])) \rceil + \lceil \sum_{j=0}^{|y|-1} (-\log(\hat{\pi}_S(y[j] \cup \{w[j]\})) + \log(\hat{\pi}_S(w[j]))) \rceil.$$

Since

$$\mathcal{I}_{\hat{\pi}_S}(u_w) = -\sum_{j=0}^{|u_w|-1} \log(\hat{\pi}_S(u_w[j])) = -\sum_{j=0}^{|y|-1} \log(\hat{\pi}_S(y[j] \cup \{w[j]\})) - \sum_{j=0}^{|z|-1} \log(\hat{\pi}_S(z[j]))$$

we have (4.5):

$$K(u_w) \leq K(w) + \mathcal{I}_{\hat{\pi}_S}(u_w) - \mathcal{I}_{\hat{\pi}_S}(w) + c.$$

□

**Proof** (of Theorem 4.5). Let  $\sigma$  and  $T$  be as given. Choose  $c_\sigma \in \mathbb{N}$  as in Theorem 3.3, and choose  $c \in \mathbb{N}$  as in Lemma 4.6. Then, for every prefix  $w \sqsubseteq \sigma * T$ , we have

$$\begin{aligned} \mathcal{I}_{\hat{\pi}_S}(u_w) - c_\sigma &\leq K(u_w) \\ &\leq \mathcal{I}_{\hat{\pi}_S}(u_w) + K(w) - |w|(\log(m-1) - \log(k-1 + A(m-k))) + c, \end{aligned}$$

so

$$\begin{aligned} K(w) &\geq |w|(\log(m-1) - \log(k-1 + A(m-k))) - c_\sigma - c \\ &= |w|(\log(m-1) - \log(k-1 + A(m-k)) - o(1)) \end{aligned}$$

as  $|w| \rightarrow \infty$ . So if  $w \rightarrow \sigma * T$  denotes  $w \sqsubseteq \sigma * T$  and  $|w| \rightarrow \infty$  we have that

$$\begin{aligned} \dim^{\pi_S}(\sigma * T) &= \liminf_{w \rightarrow \sigma * T} \frac{K(w)}{\mathcal{L}_{\pi_S}(w)} \\ &\geq \frac{\log(m-1) - \log(k-1 + A(m-k))}{-\log a}. \end{aligned}$$

□

To prove the right-hand inequality in (4.3) we need a strategy by which  $T$  can cancel as much as the randomness in  $\sigma$  as possible. A tempting strategy for this is  $T = 0^\infty$ , which always chooses the minimum element of the set chosen by  $\sigma$ . Consider this coder  $T$  in the following specific context.

**Example 4.7** Let  $F$  and  $\sigma$  be as in Example 4.2, and let  $T = 0^\infty$ . It is easy to see that the outcome  $\sigma * T$  is  $\alpha$ -random, where  $\alpha \in \Delta(\Sigma_3)$  is given by  $\alpha(0) = \frac{2}{3}$ ,  $\alpha(1) = \frac{1}{3}$ , and  $\alpha(2) = 0$ . It follows by Theorem 7.7 of [16] that  $\dim(\sigma * T) = \mathcal{H}_3(\alpha) = 1 - \frac{2}{3 \log 3} \approx 0.58$ , whence by Observation 3.1 that  $\dim(S(\sigma * T)) = (\log 3) - \frac{2}{3} \approx 0.918$ .

This example shows that the coder  $T = 0^\infty$  does indeed cancel some of the randomness in  $\sigma$ , but not enough to reach the left endpoint of the spectrum claimed in Example 4.2. The following theorem uses a nonconstructive strategy to establish the right-hand inequality of (4.3).

**Theorem 4.8** For every similarity-random selector  $\sigma$ , there is a coder  $T$  such that

$$\dim^{\pi_S}(\sigma * T) \leq \frac{\log m - \log k}{-\log A},$$

where  $A = \max\{\pi_S(i) \mid i \in \Sigma_m\}$ .

**Proof.** In this proof we use the following notation. For  $i \in \Sigma_m$  we refer to  $\hat{\pi}_S(i)$  as the sum of  $\hat{\pi}_S$  for the sets containing  $i$ , that is

$$\hat{\pi}_S(i) = \sum_{i \in B} \hat{\pi}_S(B).$$

Let  $\alpha > 1 - \log k / \log m$ . Let  $u \in \Sigma_m^*$ ,  $n \in \mathbb{N}$ . We define the set

$$Z_n^u = \{\sigma \mid \forall T \text{ with } u \sqsubseteq T, K(\sigma * T[0..n-1]) \geq \alpha n \log m\}.$$

Let  $i \in \Sigma_m$  be such that  $\pi_S(i) \geq \frac{1}{m}$ . Then  $\hat{\pi}_S(i) \geq \frac{k}{m}$ .

If  $\sigma \in Z_n^u$ , for all  $T$  extending  $u$ ,  $\sigma * T[0..n-1]$  does not end in  $(1-\alpha)n$  occurrences of the symbol  $i$  and therefore

$$\begin{aligned} \hat{\pi}_S(Z_n^u) &\leq \left(1 - (\hat{\pi}_S(i))^{(1-\alpha)n}\right)^{k^{\alpha n - |u|}} \\ &\leq \left(1 - \left(\frac{k}{m}\right)^{(1-\alpha)n}\right)^{k^{\alpha n - |u|}} \\ &\approx e^{-k^n / m^{(1-\alpha)n}} \rightarrow_n 0. \end{aligned}$$

So  $\{Z_n^u\}$  is a  $\hat{\pi}_S$ -Martin-Löf test for each  $u$ . Since  $\sigma$  is  $\hat{\pi}_S$ -random, it follows that there exist infinitely many  $n$  such that  $\sigma \notin Z_n^u$ . Therefore there is a  $T \in \Sigma_k^\infty$  such that  $K(\sigma * T[0..n-1]) < \alpha n \log m$  for infinitely many  $n$ , and  $\dim^{\pi_S}(\sigma * T) \leq \alpha \cdot \frac{\log m}{-\log A_S}$ .

Instead of a single  $\alpha$  we can take a decreasing rational sequence  $\alpha_r \rightarrow_r 1 - \log k / \log m$  and prove that there is a  $T \in \Sigma_k^\infty$  with  $\dim^{\pi_S}(\sigma * T) \leq \frac{\log m - \log k}{\log A_S}$ . □

## 5 Other Random Subfractals: upper bound for product distributions

In this section, we investigate the upper bound for points in random subfractals generated from product measures on  $\text{SEL}(\binom{m}{k})$ , i.e., measures on  $\text{SEL}(\binom{m}{k})$  that are products of measures in  $\Delta([\Sigma_m]^k)$ .

The well-known max-flow min-cut theorem is particularly useful in our investigation, so we recall the definition of flow networks for completeness.

**Definition.** A *flow network* is a tuple  $N := \langle G, s, t, c \rangle$ , where  $G = (V, E)$  is a directed graph,  $s, t \in V$ , and  $c : E \rightarrow \mathbb{R}$  is a *capacity function*. A *flow* in  $N$  is a function  $f : E \rightarrow \mathbb{R}$  that satisfies the following conditions:

1. For all  $v \in V - \{s, t\}$ ,  $\sum_{u:(u,v) \in E} f((u, v)) = \sum_{u:(v,u) \in E} f((v, u))$ .
2.  $f(e) \leq c(e)$  for all  $e \in E$ .

The *value* of a flow is defined by  $|f| = \sum_{v:(s,v) \in E} f((s, v))$ . A *cut* is a partition of  $V$  into  $S, T$  such that  $s \in S$  and  $t \in T$ . The *capacity* of a cut is defined by  $c(S, T) = \sum_{(u,v) \in S \times T} c((u, v))$ . The *maximum flow problem* is to maximize  $|f|$ . The max-flow min-cut theorem [21] states that the maximum value of a flow is equal to the minimum capacity over all cuts.

In the following, we establish conditions under which the dimensions of points in random subfractals can achieve the dimension of the fractal from which the random subfractal is generated. Our proof makes essential use of the max-flow min-cut theorem.

**Theorem 5.1** *Let  $\rho \in \Delta(\Sigma_m)$  and  $\gamma \in \Delta([\Sigma_m]^k)$  be such that the following condition holds for all  $C \subseteq \Sigma_m$*

$$\sum_{A \cap C \neq \emptyset} \gamma(A) \geq \sum_{i \in C} \rho(i). \quad (5.1)$$

*Then for every  $\gamma$ -random selector  $\sigma$  there is a  $T \in \Sigma_k^\infty$  such that  $\sigma * T$  is  $\rho$ -random.*

**Proof.** We prove this theorem by proving that there is a probability measure that depends on  $\sigma$ , according to which, a random  $T$  has the desired property. We first formulate this problem in terms of a network flow problem. We then prove that condition (5.1) implies the maximum flow in our flow network is exactly 1 and construct the desired probability measure based on a maximum flow.

Let  $G = (V, E)$  be a directed graph such that

$$V = \{s, t\} \cup [\Sigma_m]^k \cup \Sigma_m$$

and

$$E = \left\{ (s, A) \mid A \in [\Sigma_m]^k \right\} \cup \{(i, t) \mid i \in \Sigma_m\} \cup \{(A, i) \mid i \in A\}.$$

Let

$$c(e) = \begin{cases} \gamma(A) & \text{if } e = (s, A) \\ \infty & \text{if } e = (A, i) \\ \rho(i) & \text{if } e = (i, t). \end{cases}$$

be a capacity function. Then  $N = \langle G, s, t, c \rangle$  is a flow network.

Since  $\gamma$  and  $\rho$  are probability measures, it is clear that for any flow  $f : E \rightarrow \mathbb{R}$  in  $N$ ,  $|f| \leq 1$ . It is also clear that the smallest cut in  $N$  has capacity less than or equal to 1.

By the min-cut/max-flow theorem, it suffices to show that the minimum cut of  $G$  has capacity at least 1.

Note that for any cut that contains an edge in  $\{(A, i) \mid i \in A\}$ , the capacity of the cut is  $\infty$ . Any such cut cannot be a minimum cut. Let  $S = \{s\} \cup B^c \cup C^c, T = \{t\} \cup B \cup C$  be a non-trivial cut of  $G$ , where  $B \subseteq [\Sigma_m]^k$  and  $C \subseteq \Sigma_m$ . (We insist here that  $C^c \cup C = \Sigma_m$  and  $B^c \cup B = [\Sigma_m]^k$ .) Let  $B' = \{A \in [\Sigma_m]^k : A \cap C \neq \emptyset\}$ . Note that since the capacity of cut  $S, T$  is finite, no edge between  $B^c$  and  $C$  can appear in the sum of the capacity  $c(S, T)$ , therefore  $B' \subseteq B$  and it is easy to see that  $S' = \{s\} \cup B'^c \cup C^c, T' = \{t\} \cup B' \cup C$  is a cut. So  $S', T'$  has capacity at most that of  $S, T$ . The capacity of  $S', T'$  is

$$\begin{aligned} c(S', T') &= \sum_{A \in B'} c(s, A) + \sum_{i \in C^c} c(i, t) = \sum_{A \in B'} c(s, A) + 1 - \sum_{i \in C} c(i, t) \\ &= \sum_{A \cap C \neq \emptyset} \gamma(A) + 1 - \sum_{i \in C} \rho(i) \\ &\geq 1 \end{aligned} \quad \text{by condition (5.1).}$$

Therefore, the capacity of the cut  $S', T'$  is at least 1. Since  $S, T$  is an arbitrary non-trivial cut, the minimum cut capacity is 1 and the maximum flow is at least 1.

Let  $f$  be a flow of value 1 for the network  $N$ . For each  $A \in [\Sigma_m]^k$ , define the probability measure  $\nu_A \in \Delta(\Sigma_k)$  by

$$\nu_A(j) = \frac{f(A, A_j)}{\gamma(A)},$$

for each  $j \in \Sigma_k$ . ( $A_j$  is the  $j$ th element of the set  $A$  in numerical order.)

For each  $\sigma' \in \text{SEL}(\binom{m}{k})$ , define the probability measure  $\nu_{\sigma'} \in \Delta(\Sigma_k^\infty)$  by the following recursion.

$$\begin{aligned} \nu_{\sigma'}(\lambda) &= 1 \\ \nu_{\sigma'}(wa) &= \nu_{\sigma'}(w)\nu_{\sigma'(w)}(a) \end{aligned}$$

for all  $w \in \Sigma_k^*$  and all  $a \in \Sigma_k$ .

In the following, we show that for a  $\nu_\sigma$ -random  $T$ ,  $\sigma * T$  is  $\rho$ -random. We do so by showing that if  $\sigma * T$  is not  $\rho$ -random, then  $T$  is not  $\nu_\sigma$ -random relative to  $\sigma$ .

Let  $d : \Sigma_m^\infty \rightarrow \mathbb{R}$  be an optimal constructive  $\rho$ -martingale on  $\Sigma_m^\infty$  with  $d(\lambda) < 1$ .

For each  $n \in \mathbb{N}$ , let

$$Z_n = \{U \in \Sigma_m^\infty \mid (\exists u \sqsubseteq U)d(u) > 2^{2n}\}.$$

It is clear that  $\{Z_n\}_{n \in \mathbb{N}}$  is uniformly  $\Sigma_1^0$  and  $\rho(Z_n) < 2^{-2n}$ .

For each  $\sigma' \in \text{SEL}(\binom{m}{k})$  and each  $n \in \mathbb{N}$ , let

$$Y_n^{\sigma'} = \{T' \in \Sigma_k^\infty \mid (\exists u \sqsubseteq \sigma' * T')d(u) > 2^{2n}\}.$$

Let

$$X_n = \left\{ \sigma' \in \text{SEL}(\binom{m}{k}) \mid \nu_{\sigma'}(Y_n^{\sigma'}) > 2^{-n} \right\}.$$

It is clear that  $\{X_n\}_{n \in \mathbb{N}}$  is uniformly  $\Sigma_1^0$  and  $\gamma(X_n) \leq 2^{-n}$ . Since the joint distribution of  $\gamma(\sigma')$  and  $\nu_{\sigma'}$  is  $\rho$ ,

$$\gamma(X_n) \cdot 2^{-n} \leq \rho(Z_n) < 2^{-2n}.$$

Therefore,  $\gamma(X_n) < 2^{-n}$ .



Since  $\sigma$  is  $\gamma$ -random,  $|\{n \in \mathbb{N} \mid \sigma \in X_n\}| < \infty$  and thus there exists  $n_0 \in \mathbb{N}$  such that for all  $n_0 \leq n \in \mathbb{N}$ ,

$$\nu_\sigma(Y_n^\sigma) \leq 2^{-n}.$$

Note that  $\{Y_n^\sigma\}_{n \in \mathbb{N}}$  is uniformly  $\Sigma_1^0$  in  $\sigma$ . Since  $\sigma * T$  is not  $\rho$ -random,  $T \in \bigcap_{n \in \mathbb{N}} Y_n^\sigma$ , which is a  $\sigma$ -effective  $\nu_\sigma$  measure 0 set. Therefore  $T$  is not  $\nu_\sigma$ -random relative to  $\sigma$ .  $\square$

The flow network construction in the proof of Theorem 5.1 not only tells us when the points in a random subfractal can achieve the dimension of the original fractal, but also provides us a way to find out the maximum dimension of points in a random subfractal when the dimension of the original fractal is not achievable.

**Theorem 5.2** *Let  $\rho \in \Delta(\Sigma_m)$  and  $\gamma \in \Delta([\Sigma_m]^k)$  be such that condition (5.1) holds. Then for every  $\gamma$ -random selector  $\sigma$  there is a  $T \in \Sigma_k^\infty$  such that  $\dim(S(\sigma * T)) = \dim(F_\sigma(S)) = \dim_{\mathbb{H}}(F(S)) \cdot \dim^{\pi_S}(\sigma * T)$ , with*

$$\dim^{\pi_S}(\sigma * T) = \frac{\mathcal{H}(\rho)}{\mathcal{H}(\rho) + D(\rho \parallel \pi_S)},$$

which is the  $\pi_S$ -dimension of a  $\rho$ -random sequence. (Where

$$D(\rho \parallel \pi_S) = E_\rho \log \frac{\rho}{\pi_S}$$

is the Kullback-Leibler divergence between two probability measures.)

**Proof.** The result follows from Moran Theorem (equation (1.2)), Observation 3.1 and Theorem 5.1 together with the divergence formula in Theorem 3.5 of [17].  $\square$

**Corollary 5.3** *Let  $\gamma \in \Delta([\Sigma_m]^k)$  be such the following condition holds for all  $C \subseteq \Sigma_m$*

$$\sum_{A \cap C \neq \emptyset} \gamma(A) \geq \sum_{i \in C} \pi_S(i). \quad (5.2)$$

Then for every  $\gamma$ -random selector  $\sigma$  there is a  $T \in \Sigma_k^\infty$  such that  $\sigma * T$  is  $\pi_S$ -random, and

$$\dim(S(\sigma * T)) = \dim(F_\sigma(S)) = \dim_{\mathbb{H}}(F(S)).$$

**Corollary 5.4** *Let  $\gamma \in \Delta([\Sigma_m]^k)$ . Then for every  $\gamma$ -random selector  $\sigma$  there is a  $T \in \Sigma_k^\infty$  such that  $\dim(S(\sigma * T)) = \dim(F_\sigma(S)) = \dim_{\mathbb{H}}(F(S))A$ , where*

$$A = \max\{E_\rho \log \pi_S \mid \rho \text{ satisfies condition (5.1)}\}.$$

**Note.** Maximizing  $E_\rho \log \pi_S$  is equivalent to minimizing  $\mathcal{H}(\rho) + D(\rho \parallel \pi_S)$ .

**Remark.** Condition (5.1) is equivalent to requiring

$$\sum_{A \in \mathcal{C}} \gamma(A) \leq \sum_{i \in \bigcup_{A \in \mathcal{C}} A} \rho(i) \quad (5.3)$$

for all  $\mathcal{C} \subseteq [\Sigma_m]^k$ . This is because both  $\gamma$  and  $\rho$  are probability measures, i.e., both sum to 1, and the problem is symmetric. Also note that the existence of  $T \in \Sigma_k^\infty$  such that  $\sigma * T$  is  $\pi_S$ -random implies condition (5.2).

## 6 Other Random Subfractals: lower bound for general computable distribution

In the following theorem, we provide a dimension lower bound tool for more general computable probability measures on  $\text{SEL}(\binom{m}{k})$ .

**Theorem 6.1** *Let  $\gamma \in \Delta(\text{SEL}(\binom{m}{k}))$  be computable. Let  $\sigma$  be an  $\gamma$ -random selector.*

*For each  $A \in [\Sigma_m]^k$ ,  $U \in ([\Sigma_m]^k)^*$ ,  $i \in \Sigma_m$ , and  $w \in \Sigma_m^*$  define  $\rho(A|U, i)$  and  $\rho_w \in \Delta(\text{SEL}(\binom{m}{k}))$  as follows.*

$$\rho(A|U, i) = \frac{\gamma(A|U)}{\sum_{B \in [\Sigma_m]^k, i \in B} \gamma(B|U)},$$

$$\rho_w(UA) = \begin{cases} 1 & U = A = \lambda, \\ \rho_w(U)\rho(A|U, w[|T| - 1]) & (\exists T \in \Sigma_k^{\text{depth}(UA)}) UA * T \sqsubseteq w, |U * T| = |UA * T| - 1, \\ \rho_w(U)\gamma(A|U) & (\exists T \in \Sigma_k^{\text{depth}(UA)}) UA * T = U * T \sqsubseteq w, \\ 0 & \text{otherwise.} \end{cases}$$

Fix  $T \in \Sigma_k^\infty$  and let  $x = \sigma * T$ . Then

$$\dim(x) \geq \liminf_{n \rightarrow \infty} \frac{\log \left( \frac{\prod_{i \in \mathcal{U}_x[0..n-1]} \rho_{x[0..n-1]}(\sigma[i]|\sigma[0..i-1])}{\prod_{i \in \mathcal{U}_x[0..n-1]} \gamma(\sigma[i]|\sigma[0..i-1])} \right)}{n \log m},$$

where  $\mathcal{U}_w = \left\{ \text{index}^{(k)}(w') \mid w \neq w' \sqsubseteq w \right\}$  for all  $w \in \Sigma_m^*$ .

**Proof.** In here and hereafter,  $\text{depth}(U) = 1 + \max \left\{ |s_i^{(k)}| \mid i < |U| \right\}$ , is intuitively the depth (the number of layers of nodes) in the tree defined by  $U$ , as  $U$  is a prefix of some selector, which we regard as a labeled tree.

Note that when  $|T| \geq \text{depth}(U)$ ,  $T$  specifies a branch of  $U$  to as far as  $U$  allows and it is possible that only a proper prefix of  $T$  is used.

When  $UA * T = U * T$  for some  $T \in \Sigma_k^{\text{depth}(UA)}$  and  $U * T \sqsubseteq w$ , then the last symbol  $A$  in  $UA$  is not related to  $w$  and therefore the knowledge of  $w$  does not help predicting  $A$  given  $U$ .

When  $UA * T \sqsubseteq w$  and  $|U * T| = |w| - 1$ , a bit of  $w$  is partially determined by the last bit  $A$  in  $UA$  and hence the knowledge of  $w$  obviously gives some information on what  $A$  should be given  $U$ .

Define the following subprobability measure.

Let

$$\rho(U) = \sum_{w \in \Sigma_m^*} 2^{-K(w)} \rho_w(U).$$

Since  $\gamma$  is computable and the inverse of the  $*$  operator is computable,  $\rho$  is constructive and is dominated by the optimal constructive subprobability supermeasure.

Let

$$d(U) = \frac{\rho(U)}{\gamma(U)}.$$

Then  $d$  is a constructive  $r$ -supermartingale.

Since  $\sigma$  is  $\gamma$ -random,

$$\limsup_{n \rightarrow \infty} d(\sigma[0..n-1]) < \infty.$$

Let  $w \in \Sigma_m^*$ . Let  $U \in ([\Sigma_m]^k)^*$  such that there exists some  $T' \in \Sigma_k$  such that  $U * T' = w$  and  $\text{depth}(U) = |w|$  and for any  $A'$   $\text{depth}(UA') = |w| + 1$ .

Note that  $|\mathcal{U}_w| = \text{depth}(U) = |w|$  and that  $\mathcal{U}_w$  is intuitively the positions along  $U$  where the digits of  $w$  directly depend on. Let  $\mathcal{U}_w^c = \{j < |U| \mid j \notin \mathcal{U}\}$ . Then

$$\begin{aligned} d(U) &= \frac{\rho(U)}{\gamma(U)} \\ &= \frac{\sum_{w' \in \Sigma_m^*} 2^{-K(w')} \rho_{w'}(U)}{\gamma(U)} \\ &\geq 2^{-K(w)} \frac{\rho_w(U)}{\gamma(U)} \\ &= 2^{-K(w)} \frac{\prod_{i=0}^{|U|-1} \rho_w(U[i]|U[0..i-1])}{\prod_{i=0}^{|U|-1} \gamma(U[i]|U[0..i-1])} \\ &= 2^{-K(w)} \frac{\prod_{i \in \mathcal{U}_w} \rho_w(U[i]|U[0..i-1])}{\prod_{i \in \mathcal{U}_w} \gamma(U[i]|U[0..i-1])}. \end{aligned}$$

Since  $\sigma$  is  $\gamma$ -random, for every  $U \sqsubseteq \sigma$

$$d(U) \leq \mathcal{O}(1).$$

Therefore,

$$2^{-K(w)} \frac{\prod_{i \in \mathcal{U}_w} \rho_w(U[i]|U[0..i-1])}{\prod_{i \in \mathcal{U}_w} \gamma(U[i]|U[0..i-1])} \leq 1,$$

i.e.,

$$2^{K(w)} \geq \frac{\prod_{i \in \mathcal{U}_w} \rho_w(U[i]|U[0..i-1])}{\prod_{i \in \mathcal{U}_w} \gamma(U[i]|U[0..i-1])},$$

and

$$K(w) \geq \log \left( \frac{\prod_{i \in \mathcal{U}_w} \rho_w(U[i]|U[0..i-1])}{\prod_{i \in \mathcal{U}_w} \gamma(U[i]|U[0..i-1])} \right).$$

Let  $x \in \sigma * \Sigma_k^\infty$  be a sequence. So the constructive dimension of  $x$  is

$$\begin{aligned} \dim(x) &= \liminf_{n \rightarrow \infty} \frac{K(x[0..n-1])}{n \log m} \\ &\geq \liminf_{n \rightarrow \infty} \frac{\log \left( \frac{\prod_{i \in \mathcal{U}_x[0..n-1]} \rho_x[0..n-1](\sigma[i]|\sigma[0..i-1])}{\prod_{i \in \mathcal{U}_x[0..n-1]} \gamma(\sigma[i]|\sigma[0..i-1])} \right)}{n \log m}. \end{aligned}$$

□

**Remark.** In this theorem, the bound only depends on  $\gamma$ . It is easy to verify by substituting the correct probability measure that Theorem 4.5 can be derived from Theorem 6.1.

## Acknowledgment

We thank an anonymous referee and the editor for their valuable comments.

## References

- [1] K. B. Athreya, J. M. Hitchcock, J. H. Lutz, and E. Mayordomo. Effective strong dimension, algorithmic information, and computational complexity. *SIAM Journal on Computing*, 37:671–705, 2007.
- [2] G. Barmpalias, P. Brodhead, D. Cenzer, S. Dashti, and R. Weber. Algorithmic randomness of closed sets. *J. Log. Comput.*, 17(6):1041–1062, 2007.
- [3] P. Billingsley. Hausdorff dimension in probability theory ii. *Illinois Journal of Mathematics*, 5:291–298, 1961.
- [4] V. Brattka and K. Weihrauch. Computability on subsets of Euclidean space I: Closed and compact subsets. *Theoretical Computer Science*, 219:65–93, 1999.
- [5] M. Braverman and S. Cook. Computing over the reals: Foundations for scientific computing. *Notices of the AMS*, 53(3), 2006.
- [6] T. M. Cover and J. A. Thomas. *Elements of Information Theory*. John Wiley & Sons, Inc., New York, N.Y., 1991.
- [7] D. Diamondstone and B. Kjos-Hanssen. Members of random closed sets. In K. Ambos-Spies, B. Löwe, and W. Merkle, editors, *CiE*, volume 5635 of *Lecture Notes in Computer Science*, pages 144–153. Springer, 2009.
- [8] R. Dougherty, J. H. Lutz, R. D. Mauldin, and J. Teutsch. Translating the Cantor set by a random real. *Transactions of the American Mathematical Society*, 366:3027–3041, 2014.
- [9] K. Falconer. Dimensions and measures of quasi self-similar sets. *Proc. Amer. Math. Soc.*, 106:543–554, 1989.
- [10] K. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. Wiley, second edition, 2003.
- [11] A. Grzegorzcyk. Computable functionals. *Fundamenta Mathematicae*, 42:168–202, 1955.
- [12] J. M. Hitchcock. Correspondence principles for effective dimensions. *Theory of Computing Systems*, 38(5):559–571, 2005.
- [13] K.-I. Ko. *Complexity Theory of Real Functions*. Birkhäuser, Boston, 1991.
- [14] D. Lacombe. Extension de la notion de fonction recursive aux fonctions d’une ou plusieurs variables reelles, and other notes. *Comptes Rendus*, 240:2478-2480; 241:13-14, 151-153, 1250-1252, 1955.
- [15] M. Li and P. M. B. Vitányi. *An Introduction to Kolmogorov Complexity and its Applications*. Springer-Verlag, Berlin, 2008. Third Edition.
- [16] J. H. Lutz. The dimensions of individual strings and sequences. *Information and Computation*, 187:49–79, 2003.

- [17] J. H. Lutz. A divergence formula for randomness and dimension. *Theoretical Computer Science*, 412:166–177, 2011.
- [18] J. H. Lutz and E. Mayordomo. Dimensions of points in self-similar fractals. *SIAM Journal on Computing*, 38:1080–1112, 2008.
- [19] E. Mayordomo. A Kolmogorov complexity characterization of constructive Hausdorff dimension. *Information Processing Letters*, 84(1):1–3, 2002.
- [20] P. A. P. Moran. Additive functions of intervals and Hausdorff measure. *Proceedings of the Cambridge Philosophical Society*, 42:15–23, 1946.
- [21] C. H. Papadimitriou and K. Steiglitz. *Combinatorial Optimization: Algorithms and Complexity*. Dover, 1998.
- [22] M. B. Pour-El and J. I. Richards. *Computability in Analysis and Physics*. Springer-Verlag, 1989.
- [23] M. van Lambalgen. *Random Sequences*. PhD thesis, Department of Mathematics, University of Amsterdam, 1987.
- [24] K. Weihrauch. *Computable Analysis. An Introduction*. Springer-Verlag, 2000.